

RESEARCH ARTICLE

Complications of stent placement in patients with esophageal cancer: A systematic review and network meta-analysis

Amin Doosti-Irani^{1,2}, Mohammad Ali Mansournia², Abbas Rahimi-Foroushani², Peiman Haddad³, Kourosh Holakouie-Naieni^{2*}

1 Department of epidemiology, school of public health, Hamadan University of medical sciences, Hamadan, Iran, **2** Department of epidemiology and biostatistics, school of public health, Tehran University of medical sciences, Tehran, Iran, **3** Radiation Oncology Research Center, Cancer Institute, Tehran University of Medical Sciences, Tehran, Iran

* holakoik@hotmail.com



OPEN ACCESS

Citation: Doosti-Irani A, Mansournia MA, Rahimi-Foroushani A, Haddad P, Holakouie-Naieni K (2017) Complications of stent placement in patients with esophageal cancer: A systematic review and network meta-analysis. PLoS ONE 12 (10): e0184784. <https://doi.org/10.1371/journal.pone.0184784>

Editor: John Green, University Hospital Llandough, UNITED KINGDOM

Received: December 13, 2016

Accepted: August 30, 2017

Published: October 2, 2017

Copyright: © 2017 Doosti-Irani et al. This is an open access article distributed under the terms of the [Creative Commons Attribution License](https://creativecommons.org/licenses/by/4.0/), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Data Availability Statement: All relevant data are within the paper and its Supporting Information files.

Funding: This study supported by Tehran University of Medical Sciences (TUMS). We would like to thank Vic-Chancellor of Research and Technology of TUMS for financial support of this study. The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

Abstract

Background

Palliative treatments and stents are necessary for relieving dysphagia in patients with esophageal cancer. The aim of this study was to simultaneously compare available treatments in terms of complications.

Methods

Web of Science, Medline, Scopus, Cochrane Library and Embase were searched. Statistical heterogeneity was assessed using the Chi² test and was quantified by I². The results of this study were summarized in terms of Risk Ratio (RR). The random effects model was used to report the results. The rank probability for each treatment was calculated using the p-score.

Results

Out of 17855 references, 24 RCTs reported complications including treatment related death (TRD), bleeding, stent migration, aspiration, severe pain and fistula formation. In the ranking of treatments, thermal ablative therapy (p-score = 0.82), covered Evolution® stent (p-score = 0.70), brachytherapy (p-score = 0.72) and antireflux stent (p-score = 0.74) were better treatments in the network of TRD. Thermal ablative therapy (p-score = 0.86), the conventional stent (p-score = 0.62), covered Evolution® stent (p-score = 0.96) and brachytherapy (p-score = 0.82) were better treatments in the network of bleeding complications. Covered Evolution® (p-score = 0.78), uncovered (p-score = 0.88) and irradiation stents (p-score = 0.65) were better treatments in network of stent migration complications. In the network of severe pain, Conventional self-expandable nitinol alloy covered stent (p-score = 0.73), polyflex (p-score = 0.79), latex prosthesis (p-score = 0.96) and brachytherapy (p-score = 0.65) were better treatments.

Competing interests: The authors have declared that no competing interests exist.

Conclusion

According to our results, thermal ablative therapy, covered Evolution® stents, brachytherapy, and antireflux stents are associated with a lower risk of TRD. Moreover, thermal ablative therapy, conventional, covered Evolution® and brachytherapy had lower risks of bleeding. Overall, fewer complications were associated with covered Evolution® stent and brachytherapy.

Introduction

Esophageal cancer includes squamous cell carcinoma and adenocarcinoma. It has an aggressive nature, a poor prognosis and a low five-year survival rate [1]. More than 50% of patients with esophageal cancer are diagnosed with an advanced stage of the disease, for which surgery is not appropriate. Therefore, palliative treatments are necessary for relieving the associated dysphagia [2]. Certain treatment options have been developed to relieve the pain and dysphagia, including the stent, brachytherapy, and external radiotherapy [3]. Different types of stents such as Self-expandable metallic stents (SEMS), Self-expandable plastic stents (SEPS), polyflex, and antireflux, have been developed and compared in randomized control trials (RCT) [4]. However, the optimal treatment intervention has not been determined yet [3]. Up to now, many RCTs have been conducted to assess the effectiveness of stents and palliative treatments among patients with advanced esophageal cancer [5–10]. These RCTs have reported adverse events for stent placement and other palliative treatments such as brachytherapy and thermal ablative treatment in patients, including TRD, bleeding, stent migration, fistula formation, and aspiration. However, there is no consensus as to which stent has lower complications [3].

A network meta-analysis simultaneously comparing all available treatments can prove useful in the selection of better treatments [11]. Therefore, the aim of this network meta-analysis was to simultaneously compare available palliative treatments in terms of complications, including TRD, bleeding, stent migration, aspiration, severe pain and fistula formation, and to rank these treatment interventions in patients with esophageal cancer.

Methods

Search strategy and selection criteria

This network meta-analysis is a part of a comprehensive systematic review which has simultaneously compared all the treatment interventions for esophageal cancer. The protocol of this systematic review has been registered in PROSPERO (ID: CRD42015023950) and has also been published [12].

A search strategy was developed to obtain relevant RCTs on the evaluation of treatment interventions for esophageal cancer. The search strategy for this review included the following Keywords: #1: Esophageal cancer [tw]; #2: Esophageal squamous cell carcinoma [Mesh terms]; #3: Esophageal Neoplasms [Mesh terms]; #4: #1 OR #2 OR #3; #5: Randomized controlled trial [Mesh terms]; #6: Randomized clinical trial [tw]; #7: #5 OR #6; #8: Radiotherapy [Mesh terms]; #9: Stents [Mesh terms]; #10: Brachytherapy [Mesh terms]; #11: Palliative Care [Mesh terms]; #12: #8 OR #9 OR #10 OR #11; #13: #4 AND #7 AND #12.

The international databases searched until July 2017 included Web of Science, Medline, Scopus, Cochrane Library and Embase. During the hand-search, the reference lists of the included RCTs and relevant published systematic reviews and meta-analyses were also

scanned. The authors of the included studies were contacted. In addition, the following websites of relevant conferences were searched to obtain unpublished articles:

International Gastric Cancer Association; available from http://www.igca.info/news/dec2012_02.html

The International Society for Diseases of the Esophagus; available from <http://www.isde.net/events>

Cancer Research UK; available from: <http://www.cancercentre.ox.ac.uk/events/sponsored-events/symposium-on-oesophageal-cancer/>

World Organization for Specialized Studies on Diseases of the Esophagus; available from: <http://www.oeso.org/index.html>

Gastroenterology Conference Map; available from: <http://www.mdlinx.com/gastroenterology/conference-map.cfm>

All RCTs that had evaluated stent placement or palliative treatments of esophageal cancer were retrieved in this review. We put no restriction on the time, location and language of the published RCTs. The inclusion criteria for this review were, RCTs that included patients with either histology of esophageal cancer i.e. squamous cell carcinoma and/or adenocarcinoma. Cohort studies and non-randomized clinical trials were excluded.

Data extraction and quality assessment

Two investigators (ADI & MAM) screened the titles and abstracts of the retrieved RCTs independently. They reviewed the full texts of relevant RCTs to assess the eligibility for inclusion in the network meta-analysis. Any disagreement between the authors was resolved by discussion and -if need be- by the judgment of other authors. The following data were extracted using a predefined data sheet: year of publication, location, duration of study (months), stage of cancer, type of palliative treatment in each arm of the RCT in detail, number of randomized patients in each arm of the RCT, numbers of males and females, mean/median age of participants and the numbers of complications among patients in each arm of the included RCT. The risk of bias was assessed using Cochrane's tools [13]. The details of used items from Cochrane's tools in this review were reported in the protocol [12].

Outcome

The outcomes of interest in this study were TRD, bleeding, stent migration, aspiration, severe pain and fistula formation among patients with esophageal cancer who had received palliative treatment interventions.

Statistical analysis


In the first stage, the network of treatment interventions was drawn using the *netgraph* command in R software. We performed a pairwise meta-analysis for each comparison using the random effect model. A Risk Ratio with 95% CI was calculated for each complication in two by two comparisons. The statistical heterogeneity was assessed using the Chi² test and the heterogeneity across each comparison was quantified using I² statistics [14]. The similarity assumption was assessed clinically and epidemiologically in terms of effect modifiers in the arms of the RCTs. The consistency assumption was assessed using loop-specific and design by treatment interaction methods [15]. The publication bias for each complication was assessed visually by the adjusted network funnel plot using Stata 13 (Stata Corp, College Station, TX, USA) [16]. The results of this random effect network meta-analysis were summarized in terms of Risk Ratio with 95% Confidence Interval (CI).

The rank probabilities of the treatments were calculated using the p-score. The p-score for treatment i is one minus the one-sided p-value $p[j]$ of accepting the alternative hypothesis. Therefore, the p-score for each treatment is the mean of all $1-p[j]$. The P-score is a value between zero and one, where a larger value indicates a better treatment [17]. The statistical analysis was performed using R, version 3.3.1 (2016-06-21), with the netmeta package for network meta-analysis.

Results

Overall, 17855 references were retrieved after removing duplicate references. After screening the titles and abstracts, 832 RCTs' full texts remained for review. Upon checking the eligibility criteria, 24 RCTs [5, 7–10, 18–36] had reported the complications of stents and palliative treatments, so they were included in our network meta-analysis (Fig 1 and Table 1).

As shown in Figs 2 and 3, and S1, S3, S5 and S7 Figs, there were no closed loops in the networks, so we could not assess the consistency assumption of the complications. However, we assessed the heterogeneity in the two by two comparisons and the entire networks. The results showed no significant heterogeneity in the networks in either of the complications (S1–S6 Tables). Based on the adjusted funnel plot there was no evidence of publication bias for the set of studies related to each complication (S9–S14 Figs).



From: Moher D, Liberati A, Tetzlaff J, Altman DG, The PRISMA Group (2009). Preferred Reporting Items for Systematic Reviews and Meta-Analyses: The PRISMA Statement. PLoS Med 6(7): e1000097. doi:10.1371/journal.pmed1000097

For more information, visit www.prisma-statement.org.

Fig 1. A flow chart depicting the stages of retrieving articles and checking the eligibility criteria for network meta-analysis.

<https://doi.org/10.1371/journal.pone.0184784.g001>

Table 1. Characteristics of included randomized control trials.

Author	Quality	Country	Treatments	n1	n2	proportion of male in arm 1	Proportion of male in arm 2	Mean age 1	Mean age 2	TRD 1	TRD 2	Bleeding 1	Bleeding 2	Stent migration 1	Stent migration 2	Aspiration 1	Aspiration 2	Sever pain 1	Sever pain 2	Fistula 1	Fistula 2
Amdal (2013)	I	Norway	T1:SEMS +BT T2: BT	21	20	0.62	0.65	74	73			1	0	1	0	3	0				
Conio (2007)	I	International	T1: Polyflex T2: Ultraflex stent	47	54	0.83	0.83	74.9	69.1	2	4	2	0	7	1					1	2
Dallal (2001)	I	UK	T1: Thermal ablative therapy T2: Metallic stent	34	31	0.53	0.52	75	77	1	2	0	3	0	1					3	0
De Palma (1996)	L	Italy	T1: Plastic stent T2: Metallic stent	20	19	0.75	0.84	69.4	67.8	3	0	1	0								
Guo (2008)	H	China	T1: Irradiation stent T2: Conventional stent	27	26	0.70	0.77	72.2	69.5	6	5	9	7	2	3	1	2	8	7	1	0
Homs (2004)	I	Netherlands	T1: BT T2: SEMS 18	101	108	0.75	0.80	69	69	1	6	5	14	3*	18	1	1	1	3	3	3
Homs (2004)	I	Netherlands	T1: FerX-Ella + antireflux T2: FerX-Ella	15	15	0.80	0.80	68	69	0	0	2	1	5	2	0	1	1	1		
Javed (2012)	H	India	T1: Ultraflex stent T2: Ultraflex stent+RT	37	42	0.73	0.69	58.1	58.6	0	0			9	6						
Knyrim (1993)	L	Germany	T1: Plastic stent T2: Metallic stent	21	21	0.00	0.00	66.8	64.8	6	3			5	0	1	0			1	2
O'Donnell (2002)	L	UK	T1: Plastic stent T2: Metallic stent	25	25	0.00	0.00	72.3	72.9	5	6			3	2						
Power (2007)	H	Ireland	T1: Conventional stent T2: Antireflux stent	25	24	0.68	0.58	73.9	64.8			0	0	0	2			1	2		
Sabharwal (2008)	I	UK	T1: Antireflux stent T2: Ultraflex stent + omeprazole	22	26	0.68	0.81	71.4	66.3	0	1	1	2	7	6			2	9	0	1
Sabharwal (2003)	I	UK	T1: Ultraflex stent T2: Flamingo stent	31	22	0.81	0.68	71.6	61.2	5	4	1	1	2	1						
Shenfine (2009)	H	UK	T1: Rigid plastic stent T2: Non-Stent	57	50	0.70	0.78	78.2	76.9			12	6	17	8	2	0	8	7		

(Continued)

Table 1. (Continued)

Author	Quality	Country	Treatments	n1	n2	proportion of male in arm 1	Proportion of male in arm 2	Mean age 1	Mean age 2	TRD 1	TRD 2	Bleeding 1	Bleeding 2	Stent migration 1	Stent migration 2	Aspiration 1	Aspiration 2	Sever pain 1	Sever pain 2	Fistula 1	Fistula 2
Vakli (2001)	I	Italy	T1: Metallic stent T2: Uncovered stent	32	30	0.00	0.00	74	71			8	2	4	2			12	15		
van Heel (2012)	L	Netherlands	T1: Ultraflex stent T2: Covered Evolution stent	40	40	0.80	0.65	67	67	10	7	7	0	3	1	3	2	5	6		
Wenger (2006)	I	Sweden	T1: Antireflux T2: Conventional	19	22	0.68	0.59	75	73	0	0	1	1	2	3						
Wenger (2005)	H	Sweden	T1: SEMS T2: BT	30	30	0.60	0.70	74	72	2	2	1	0	2	-					2	1
White (2015)	H	Kenia	T1: SEMS 18 T2: SEMS 23	50	50	0.62	0.58	61.8	57.1					3	0			1	0		
Zhu (2014)	H	China	T1: Irradiation Stent T2: CSENAACS	80	80	0.76	0.66	71	71			5	5	0	0	11	14	17	15	6	5
Laasch (2002)	L	UK	T1: Open Stent T2: Antireflux stent	25	25	0.76	0.60	69	69	1	0			3	4			1	0		
Siersema (1998)	I	Netherlands	T1: Latex prosthesis T2: Metallic stent	38	37	0.76	0.70	65.2	67.6	4	1	5	3	3	0			6	12	1	0
Siersema (2001)	L	Netherlands	T1: Ultraflex stent T2: Flamingo stent	34	33	0.82	0.67	67.7	71.4	1	0	5	3	6	3			3	6		
Verschuur (2008)	I	Italy	T1: Ultraflex stent T2: Polyflex stent	42	41	0.67	0.68	69	70	1	2	5	5	7	12	1	0	1	0	2	0

H: high quality; I: intermediate quality; L: low quality; T1: treatment 1, T2: treatment 2; BT: Brachytherapy; n1: sample size in arm 1, n2: sample size in arm 2; CSENAACS: Conventional self-expandable nitinol alloy covered stent; SEMS: Self-expandable metallic stents.

*In this RCT some of randomized patients in BT group received SEMS 18.

<https://doi.org/10.1371/journal.pone.0184784.t001>

Treatment-related death complication

TRD was reported in 16 RCTs, which included 1075 patients with esophageal cancer. The networks of eligible comparisons for TRD and bleeding are shown in Figs 2 and 3. The comparisons of treatments for TRD involved four independent sub-networks. In the networks, A and B, C and D, the metallic stent, the Ultraflex stent, brachytherapy and antireflux stents were reference treatments, respectively (Fig 2). According to the results of the test for heterogeneity, the I^2 statistic for network A was 15.2%, and for network B, C, and D was zero (S1 Table).

The comparison of palliative treatments with reference treatments in each network for TRD is shown in Fig 3. In network A, compared with the metallic stent, the latex prosthesis increased the risk of TRD. The relative risk (RR) was 3.89 (95% CI: 0.42, 36.33). The RR for thermal ablative therapy compared with the metallic stent was 0.46 (95% CI: 0.04, 5.19). In network B, covered Evolution® compared with Ultraflex stent decreased the risk of TRD, RR = 0.70 (95% CI: 0.30, 1.66). In network C, SEMS 18 compared to brachytherapy increased the risk of TRD, RR = 5.61, (95% CI: 0.69, 45.80). In network D, both the open stent and ‘Ultraflex plus omeprazole’ compared to antireflux stent increased the risk of TRD (Fig 3).

Results of simultaneous direct and indirect comparisons of treatments for TRD are shown in S1 Table.

In terms of ranking of treatments, thermal ablative therapy (p-score = 0.82), covered Evolution® (p-score = 0.70), brachytherapy (p-score = 0.72) and antireflux stent (p-score = 0.74) were the best treatments in networks A, B, C and D, respectively (Table 2).




Fig 2. Network of stent interventions for palliative treatments that had reported TRD in esophageal cancer; A: the metallic stent is a reference treatment, B: Ultraflex is a reference, C: Brachytherapy is a reference, and D: Antireflux is a reference.

<https://doi.org/10.1371/journal.pone.0184784.g002>




Fig 3. Forest plots for TRD in networks A, B, C, & D. Metallic stent, Ultraflex, Brachytherapy, and Antireflux are reference treatments in networks A, B, C & D, respectively.

<https://doi.org/10.1371/journal.pone.0184784.g003>

Bleeding

Bleeding was reported in 18 RCTs, which included 1374 patients with esophageal cancer.

Other characteristics of the included RCTs are shown in [Table 1](#).

In the network of bleeding complications, the metallic stent, conventional stent, Ultraflex stent, and brachytherapy were reference treatments in networks A, B, C, and D, respectively ([Fig 4](#)). Based on the results of the test for heterogeneity, the I^2 statistic for network A, B, C, and D was zero ([S2 Table](#)).

In network A, the latex prosthesis and plastic stent increased the risk of bleeding when compared to the metallic stent. The RR for latex prosthesis was 1.62 (95% CI: 0.42, 6.31) and was 2.85 (95% CI: 0.12, 65.93) for the plastic stent. On the other hand, thermal ablative therapy (RR = 0.13, 95% CI: 0.01, 2.43) and uncovered stent (RR = 0.27, 95% CI: 0.06, 1.16) decreased the risk of bleeding when compared to the metallic stent. In network B, the irradiation stent and CSENACS increased the risk of bleeding when compared to the conventional stent. In network C, the covered Evolution® stent decreased the risk of bleeding (RR = 0.07, 95% CI: 0.00, 1.13) when compared to Ultraflex. In network D, SEMS and SEMS+BT increased the risk of bleeding when compared to brachytherapy ([Fig 5](#)).

The results of simultaneous direct and indirect comparisons of treatments for bleeding are shown in [S2 Table](#).

In terms of ranking treatments for the lower risk of bleeding, thermal ablative therapy (p-score = 0.86), the conventional stent (p-score = 0.62), covered Evolution® (p-score = 0.96) and brachytherapy (p-score = 0.82) were the best treatments in networks A, B, C, and D, respectively ([Table 2](#)).

Stent migration

Stent migration was reported in 23 RCTs. However, in 4 of the RCTs patients had received BT or thermal ablative therapy [[5](#), [8](#), [19](#), [34](#)] in one arm. Since stent migration cannot be a complication of BT and thermal ablative therapy, these RCTs were excluded from the network meta-analysis. Therefore, 19 RCTs involving 1207 patients were included. The available treatments for stent migration involved three dependent networks ([S1 Fig](#)). In the network A, and B the I^2 statistic was zero, and in the network C the I^2 was 16.1% ([S3 Table](#)).

In network A, when compared to the Ultraflex stent, the polyflex stent increased the risk of stent migration 2.07 times (95% CI: 1.01, 4.67). The risk of stent migration for covered Evolution® stent, Flamingo stent, and Ultraflex stent plus radiotherapy was lower than the ultraflex

Table 2. Ranking of palliative treatments in terms of lower risk of TRD and bleeding, stent migration, aspiration, severe pain and fistula in patients with esophageal cancer.

TRD							
Network A		Network B		Network C		Network D	
Treatment	P-score	Treatment	P-score	Treatment	P-score	Treatment	P-score
Thermal ablative therapy	0.82	Covered evolution	0.70	Brachytherapy	0.72	Antireflux	0.74
Metallic stent	0.65	Polyflex	0.49	SEMS	0.69	Ultraflex +omeprazole	0.40
Plastic stent	0.38	Flamingo	0.44	SEMS18	0.09	Open stent	0.36
Latex prosthesis	0.15	Ultraflex	0.37				
Bleeding							
Network A		Network B		Network C		Network D	
Treatment	P-score	Treatment	P-score	Treatment	P-score	Treatment	P-score
Thermal ablative therapy	0.86	Conventional	0.62	Covered evolution	0.96	Brachytherapy	0.82
Uncovered stent	0.79	Antireflux	0.54	Flamingo	0.50	SEMSBT	0.41
Metallic stent	0.41	CSENACS	0.49	Ultraflex	0.34	SEMS	0.40
Latex	0.24	Irradiation stent	0.47	Polyflex	0.20	SEMS18	0.36
Plastic stent	0.20	Ultraflex + Omeprazole	0.37	-	-	-	-
Stent migration							
Network A		Network B		Network C			
Treatment	P-score	Treatment	P-score	Treatment	P-score		
Covered Evolution stent	0.78	Uncovered stent	0.88	Irradiation stent	0.65		
Flamingo stent	0.67	Metallic stent	0.69	Ultraflex stent+ omeprazole	0.52		
Ultraflex stent + RT	0.66	Plastic stent	0.27	Open stent	0.50		
Ultraflex stent	0.36	Latex prosthesis stent	0.16	Conventional stent	0.49		
Polyflex stent	0.03			Antireflux stent	0.34		
Aspiration							
Network A		Network B		Network C			
Treatment	P-score	Treatment	P-score	Treatment	P-score		
Polyflex stent	0.69	Irradiation stent	0.74	BT	0.69		
Covered evolution	0.52	CSENACS	0.46	SEMS18	0.68		
Ultraflex stent	0.29	Conventional stent	0.31	SEMSBT	0.13		
Severe pain							
Network A		Network B		Network C		Network D	
Treatment	P-score	Treatment	P-score	Treatment	P-score	Treatment	P-score
CSENACS	0.73	Polyflex stent	0.79	Latex prosthesis	0.96	BT	0.65
Conventional stent	0.71	Ultraflex	0.58	Metallic stent	0.44	SEMS23	0.63
Irradiation stent	0.63	Covered evolution	0.45	Uncovered stent	0.10	SEMS18	0.22
Antireflux	0.52	Flamingo	0.19	-	-	-	-
Open stent	0.27	-	-	-	-	-	-
Ultraflex + Omeprazole	0.14	-	-	-	-	-	-
Fistula							
Network A		Network B		Network C			
Treatment	p-score	Treatment	p-score	Treatment	p-score		
Plastic stent	0.81	Conventional stent	0.72	SEMS18	0.62		
Metallic stent	0.64	CSENACS	0.46	BT	0.59		
Latex	0.36	Irradiation	0.32	SEMS	0.29		
Thermal ablative therapy	0.19	-	-	-	-		

<https://doi.org/10.1371/journal.pone.0184784.t002>




Fig 4. The network of stent interventions for palliative treatments that reported the bleeding complication in esophageal cancer; A: metallic stent is a reference treatment, B: Conventional is a reference, C: Ultraflex is a reference, and D: Brachytherapy is a reference treatment. CSENACS: Conventional self-expandable nitinol alloy covered stent.

<https://doi.org/10.1371/journal.pone.0184784.g004>

stent; however, the 95% CIs involved the null values. In network B, the risk ratio for the latex prosthesis and plastic stents compared to the metallic stent was 6.82 (95% CI: 0.36, 127.54) and 2.87 (95% CI: 0.87, 10.64), respectively. In network C, there were no considerable differences between the conventional, Irradiation, open and ultraflex stents plus omeprazole and the




Fig 5. Forest plots for bleeding complication in networks A, B, C & D. Metallic stent, Conventional stent, Ultraflex, and Brachytherapy are reference treatments in networks A, B, C & D, respectively.

<https://doi.org/10.1371/journal.pone.0184784.g005>

Antireflux stent (S2 Fig). The results of simultaneous direct and indirect comparisons of treatments for stent migration are shown in S3 Table.

In terms of ranking, the covered Evolution® (p-score = 0.78), Uncovered (p-score = 0.88), and Irradiation stents (p-score = 0.65) were the best treatments in networks A, B and C, respectively (Table 2).

Aspiration

Nine RCTs reported aspiration as a complication of stent placement. These RCTs involved 805 esophageal cancer patients. Three RCTs involved independent comparisons, so they were excluded from the network meta-analysis [20, 21, 27]. These studies have been described in Table 1. The remaining RCTs involved three independent networks (S3 Fig). The I^2 statistic for all networks of this complication was zero (S4 Table). Forest plots drawn to compare the treatments with reference treatments in each network are shown in S4 Fig. Results of simultaneous direct and indirect comparisons of treatments for aspiration are shown in S2 Table. In terms of ranking, the Polyflex stent (p-score = 0.69), Irradiation stent (p-score = 0.74) and BT (p-score = 0.69) were the better treatments in networks A, B and C (Table 2).

Severe pain

Severe pain was reported in 14 RCTs. The CSENACS (p-score = 0.73), Polyflex stent (p-score = 0.79), Latex prosthesis (p-score = 0.96) and BT (p-score = 0.65) were better treatments in terms of lower risk of severe pain among patients in networks A, B, C and D, respectively. The network and forest plots are shown in S5 and S6 Figs. According to the results of test for heterogeneity, The I^2 statistic for network A, B, C, and D was zero (S5 Table). In addition, the results of direct and indirect comparisons are presented in S2 Table.

Fistula formation

Fistula formation was reported in 10 RCTs. The networks of interventions and forest plots drawn to compare the treatments with reference treatments in each network are shown in S7 and S8 Figs, respectively. The I^2 statistic for all networks of this complication was zero (S6 Table). The Plastic stent (p-score = 0.81), Conventional stent (p-score = 0.72), and SEMS 18 (p-score = 0.62) were better treatments in terms of lower risk of fistula formation in networks A, B, and C, respectively (Table 2).

Discussion

In this network meta-analysis, we compared the complications of palliative treatments including stents in patients with esophageal cancer. The treatments were ranked based on their lower risk of complications including TRD, bleeding, stent migration, aspiration, severe pain and fistula formation.

Based on our results, the TRD complication involved four networks. In network A, the latex prosthesis stent increased the risk of TRD 3.89 times when compared to the metallic stent. Thermal ablative therapy decreased the risk of TRD. In network B, there was no considerable difference between the covered Evolution®, Flamingo, and Ultraflex stents compared with the Ultraflex stent. In network C, when compared to brachytherapy, SEMS 18 increased the risk of TRD. In network D, the open stent and Ultraflex plus omeprazole versus antireflux stent increased the risk of TRD among patients with esophageal cancer.

Upon ranking treatments in terms of lower risk of TRD, thermal ablative therapy, covered Evolution®, brachytherapy and antireflux were better treatment interventions in the TRD

networks, respectively. However, in network C, there was no considerable difference in the p-score of brachytherapy (p-score = 0.72) and SEMS (p-score = 0.69).

In terms of lower risk of bleeding, Thermal ablative therapy, Conventional stent, Covered Evolution® stent, and Brachytherapy were ranked as best treatments in networks A, B, C, and D, respectively. The results of bleeding complication were similar to those of TRD. Based on the lower risk of stent migration, the covered Evolution® stent, uncovered stent and irradiation stent were ranked as best treatments in the networks of this complication.

The risk of aspiration was lower for the polyflex stent, Irradiation stent, and brachytherapy in the networks of this complication. However, there was no considerable difference between the first (brachytherapy) and second (SEMS 18) ranked treatments in network C. The risk of severe pain in the CSENACS, polyflex stent, latex prosthesis and brachytherapy was lower than in the other treatments in the networks.

In terms of fistula formation, the plastic stent, conventional stent and SEMS 18 were better treatments in the networks of this complication. However, it should be noted the plastic stents compared with other stents, increased the risk of TRD, bleeding, stent migration and aspiration. Therefore, we could not recommend plastic stents as a better treatment, just because the risk of fistula was lower for this stent. In addition, a recently published clinical guideline recommends against for placement of plastic stents for palliation of esophageal strictures [37].

Based on the results of a traditional meta-analysis, the risk of TRD in palliative locoregional modalities was lower than in the metallic stent (RR = 0.58, 95% CI: 0.17, 1.99). Moreover, the odds ratio (OR) for bleeding was 0.54 (95% CI: 0.29, 1.00) compared to the metallic stents [38]. These results are in line with ours.

In the aforementioned meta-analysis [38], the odds ratio of stent migration in patients who had received ultraflex stents versus other types of stents was 1.17 (95% CI: 0.71, 1.93). In our study, the risk of stent migration was higher for ultraflex stent versus covered Evolution® stent, Flamingo stent, and Ultraflex stent plus radiotherapy. However, the risk of stent migration was higher in the polyflex stent than in the ultraflex stent (RR = 2.17(1.01, 4.67)). Moreover, in our study, the risk of stent migration for the flamingo stent was lower than in the polyflex and ultraflex stents, a finding that is consistent with the aforementioned traditional meta-analysis.

According to this meta-analysis [38], the odds ratio of severe pain for the ultraflex stent versus the other stents was 0.52; 95% CI (0.19, 1.45). This result is in line with our study regarding the comparison of ultraflex versus covered Evolution® stent and Flamingo stent. However, the risk of severe pain was higher than in the polyflex stent.

The SEMS has been introduced as a selective palliative treatment in patients with esophageal cancer. The TRD in patients in which this stent had been used was reported between 0 to 1.4% [4]. In our study, the SEMS and SEMS 18 were compared with brachytherapy in network C (for TRD) and in network D (for the bleeding complication). When compared with brachytherapy, SEMS 18 increased the risk of TRD and bleeding 5.61 and 2.62 times, respectively. Likewise, compared with brachytherapy, SEMS increased the risk of bleeding by 3 times. These findings confirm the available evidence regarding the role of brachytherapy in the palliative treatment of patients with advanced stages of esophageal cancer [39]. In addition, according to the results of a recently published meta-analysis brachytherapy was recommended as a relatively safe and highly effective treatment for palliation of dysphagia [40] that is in the line of our results in terms of lower risk of TRD and bleeding for this treatment. However, despite the available evidence regarding the safety of this treatment and strong recommendation of the international guidelines [37] for use of this treatment, the lack of experience of radiation oncologist lead to the underuse of brachytherapy [41].

According to the results of one RCT, the antireflux stent had no additional advantages over the open stent [42]. But in our study, the risk of TRD in patients who had received the open stent was more than in those who had received the antireflux stent. Moreover, the p-score for the antireflux and open stents were 0.74 and 0.36, respectively.

In our network meta-analysis, the confidence intervals for RRs involved the null value in some of the networks. The reason may be due to the low number of RCTs in each network. However, the point estimates of RR in the networks of complications were considerable.

The statistical heterogeneity among the networks of complications was low. However, the statistical power of the Cochrane test is low when the number of studies included is low [3]. So these results may have been affected by the low number of studies in the networks.

According to current literature, there is controversy over the selection of the ideal stent [4]. To our knowledge, our study is the first network meta-analysis to compare the available stents in terms of six major complications (treatment-related death, bleeding, stent migration, aspiration, severe pain and fistula formation) among patients with esophageal cancer. We have provided useful evidence on the ranking of stents with lower risks of the aforementioned complications. In addition, the advantage of our study -as a network meta-analysis- over the traditional meta-analysis is the simultaneous comparison of all available treatments in each network, while traditional meta-analysis compares the stents two by two or one stent versus another [38].

There are certain limitations in this study. Firstly, palliative treatments and stent interventions involved four networks of TRD, bleeding, and severe pain, and three networks for stent migration, aspiration, and fistula formation. Therefore, we could not compare all treatment interventions simultaneously for each complication in one network. Secondly, the number of included RCTs in each network was low, so the power of networks for estimating the indirect comparison was low and the confidence intervals for indirect estimates were wide [43], and this issue can be lead to spars-data bias [44]. Therefore, more RCTs are required for future network meta-analysis.

Secondly, because of the small number of RCTs in each network and consequently low power and lack of validity of statistical tests for publication bias [45, 46], we could not be assessed the publication bias in this network meta-analysis.

We recommend that future RCTs report 'all' the complications of stent placement and other palliative treatments in patients with esophageal cancer. Thus, future systematic reviews and network meta-analyses will be able to compare stents in terms of all the major complications simultaneously.

Conclusion

Overall, the results of this network meta-analysis showed that thermal ablative therapy, covered Evolution® stents, brachytherapy and antireflux stents are associated with a lower risk of TRD. In terms of lower risk of bleeding, thermal ablative therapy, conventional stent, covered Evolution® stent and brachytherapy were better palliative treatments for patients with esophageal cancer. Based on the lower risk of stent migration, the covered Evolution®, uncovered, and Irradiation stents were better treatments. In terms of lower risk of severe pain as another major complication the CSENACS, polyflex stent, latex prosthesis and brachytherapy were better treatments.

Supporting information

S1 File. PRISMA checklist.
(DOCX)

S2 File. Search strategy for Medline.

(DOCX)

S1 Fig. Network of stent interventions for palliative treatments that reported the stent migration in esophageal cancer; A: Ultraflex stent is reference treatment, B: Metallic stent is reference, and C: Antireflux stent is reference.

(TIF)

S2 Fig. Forest plots for stent migration in network A, B, C; Ultraflex stent, Metallic stent, Anti-reflux stent are reference treatments in network A, B, and C respectively.

(TIF)

S3 Fig. Network of stent interventions for palliative treatments that reported the aspiration in patients; A: Ultraflex stent is reference treatment, B: Irradiation stent is reference, and C: Brachytherapy is reference.

(TIF)

S4 Fig. Forest plots for aspiration in network A, B, C; covered evolution stent, Irradiation stent, brachytherapy are reference treatments in network A, B, and C respectively.

(TIF)

S5 Fig. Network of stent interventions for palliative treatments that reported the severe pain in patients; Irradiation stent, Polyflex stent, Metallic stent and SEMS 18 are reference treatments in networks A, B, C and D respectively.

(TIF)

S6 Fig. Forest plots for severe pain in network A, B, C and D; Antireflux stent, Ultraflex stent, Metallic stent, and brachytherapy are reference treatments in network A, B, C and D respectively.

(TIF)

S7 Fig. Network of stent interventions for palliative treatments that reported the fistula in patients; Irradiation stent, Polyflex stent, Metallic stent and SEMS 18 are reference treatments in networks A, B, C and D respectively.

(TIF)

S8 Fig. Forest plots for fistula in network A, B, and C. Metallic stent, Irradiation stent, and brachytherapy are reference treatments in network A, B, and C respectively.

(TIF)

S9 Fig. Adjusted network funnel plot for network meta-analysis of TRD.

(TIF)

S10 Fig. Adjusted network funnel plot for network meta-analysis of bleeding complication.

(TIF)

S11 Fig. Adjusted network funnel plot for network meta-analysis of stent migration complication.

(TIF)

S12 Fig. Adjusted network funnel plot for network meta-analysis of aspiration complication.

(TIF)

S13 Fig. Adjusted network funnel plot for network meta-analysis of severe pain complication.

(TIF)

S14 Fig. Adjusted network funnel plot for network meta-analysis of fistula complication.

(TIF)

S1 Table. Simultaneous comparisons of palliative treatments using relative risk (95% CI) in terms of treatment related death among esophageal cancer patients.

(DOCX)

S2 Table. Simultaneous comparisons of palliative treatments using relative risk (95% CI) in terms of bleeding complications among esophageal cancer patients.

(DOCX)

S3 Table. Simultaneous comparisons of palliative treatments using relative risk (95% CI) in terms of stent migration among esophageal cancer patients.

(DOCX)

S4 Table. Simultaneous comparisons of palliative treatments using relative risk (95% CI) in terms of aspiration among esophageal cancer patients.

(DOCX)

S5 Table. Simultaneous comparisons of palliative treatments using relative risk (95% CI) in terms of severe pain among esophageal cancer patients.

(DOCX)

S6 Table. Simultaneous comparisons of palliative treatments using relative risk (95% CI) in terms of fistula among esophageal cancer patients.

(DOCX)

Acknowledgments

This study supported by Tehran University of Medical Sciences (TUMS). We would like to thank Vic-Chancellor of Research and Technology of TUMS for financial support of this study.

Author Contributions

Conceptualization: Amin Doosti-Irani, Peiman Haddad, Kourosh Holakouie-Naieni.

Data curation: Amin Doosti-Irani.

Formal analysis: Amin Doosti-Irani, Mohammad Ali Mansournia, Abbas Rahimi-Foroushani.

Investigation: Amin Doosti-Irani, Peiman Haddad, Kourosh Holakouie-Naieni.

Methodology: Amin Doosti-Irani, Mohammad Ali Mansournia, Abbas Rahimi-Foroushani, Kourosh Holakouie-Naieni.

Project administration: Kourosh Holakouie-Naieni.

Software: Amin Doosti-Irani, Abbas Rahimi-Foroushani.

Supervision: Kourosh Holakouie-Naieni.

Validation: Amin Doosti-Irani, Mohammad Ali Mansournia, Peiman Haddad, Kourosh Holakouie-Naieni.

Visualization: Amin Doosti-Irani.

Writing – original draft: Amin Doosti-Irani, Mohammad Ali Mansournia, Abbas Rahimi-Foroushani, Peiman Haddad, Kourosh Holakouie-Naieni.

Writing – review & editing: Amin Doosti-Irani, Mohammad Ali Mansournia, Abbas Rahimi-Foroushani, Peiman Haddad, Kourosh Holakouie-Naieni.

References

1. Ferlay J, Shin HR, Bray F, Forman D, Mathers C, Parkin DM. Estimates of worldwide burden of cancer in 2008: GLOBOCAN 2008. *International journal of cancer*. 2010; 127(12):2893–917. <https://doi.org/10.1002/ijc.25516> PMID: 21351269
2. Tytgat G, Bartelink H, Bernards R, Giaccone G, Van Lanschot J, Offerhaus G, et al. Cancer of the esophagus and gastric cardia: recent advances. *Diseases of the Esophagus*. 2004; 17(1):10–26. <https://doi.org/10.1111/j.1442-2050.2004.00371.x> PMID: 15209736
3. Sreedharan A, Harris K, Crellin A, Forman D, Everett SM. Interventions for dysphagia in oesophageal cancer. *The Cochrane database of systematic reviews*. 2009;(4):Cd005048. Epub 2009/10/13. <https://doi.org/10.1002/14651858.CD005048.pub2> PMID: 19821338.
4. Irani S, Kozarek R. Esophageal stents: past, present, and future. *Techniques in Gastrointestinal Endoscopy*. 2010; 12(4):178–90.
5. Amdal CD, Jacobsen AB, Sandstad B, Warloe T, Bjordal K. Palliative brachytherapy with or without primary stent placement in patients with oesophageal cancer, a randomised phase III trial. *Radiotherapy and Oncology*. 2013; 107(3):428–33. <https://doi.org/10.1016/j.radonc.2013.04.008> PMID: 23647761
6. Bergquist H, Wenger U, Johnsson E, Nyman J, Ejnell H, Hammerlid E, et al. Stent insertion or endoluminal brachytherapy as palliation of patients with advanced cancer of the esophagus and gastroesophageal junction. Results of a randomized, controlled clinical trial. *Diseases of the Esophagus*. 2005; 18(3):131–9. <https://doi.org/10.1111/j.1442-2050.2005.00467.x> PMID: 16045572
7. Conio MMD, Repici AMD, Battaglia GMD, De Pretis GMD, Ghezzi LMD, Bittinger MMD, et al. A Randomized Prospective Comparison of Self-Expandable Plastic Stents and Partially Covered Self-Expandable Metal Stents in the Palliation of Malignant Esophageal Dysphagia. *American Journal of Gastroenterology*. 2007; 102(12):2667–77. <https://doi.org/10.1111/j.1572-0241.2007.01565.x> PMID: 18042102
8. Dallal HJ, Smith GD, Grieve DC, Ghosh S, Penman ID, Palmer KR. A randomized trial of thermal ablation therapy versus expandable metal stents in the palliative treatment of patients with esophageal carcinoma. *Gastrointestinal Endoscopy*. 2001; 54(5):549–57. PMID: 11677469.
9. Guo JH, Teng GJ, Zhu GY, He SC, Fang W, Deng G, et al. Self-expandable esophageal stent loaded with 125I seeds: initial experience in patients with advanced esophageal cancer. *Radiology*. 2008; 247(2):574–81. Epub 2008/03/20. <https://doi.org/10.1148/radiol.2472070999> PMID: 18349316.
10. Javed A, Pal S, Dash NR, Ahuja V, Mohanti BK, Vishnubhatla S, et al. Palliative stenting with or without radiotherapy for inoperable esophageal carcinoma: A randomized trial. *Journal of Gastrointestinal Cancer*. 2012; 43(1):63–9. <https://doi.org/10.1007/s12029-010-9206-4> PMID: 20835926
11. Caldwell DM, Ades A, Higgins J. Simultaneous comparison of multiple treatments: combining direct and indirect evidence. *British Medical Journal*. 2005; 7521:897.
12. Doosti-Irani A, Mansournia MA, Rahimi-Foroushani A, Cheraghi Z, Holakouie-Naieni K. Simultaneous Comparison of Efficacy and Adverse Events of Interventions for Patients with Esophageal Cancer: Protocol for a Systematic Review and Bayesian Network Meta-analysis. *Asian Pacific journal of cancer prevention: APJCP*. 2016; 17(2):867–72. Epub 2016/03/02. PMID: 26925694.
13. Higgins JP, Green S. *Cochrane Handbook for Systematic Reviews of Interventions Version 5.1.0* [updated March 2011]; The Cochrane Collaboration; 2011. Available from: <http://handbook.cochrane.org/>.
14. Higgins JPT, Thompson SG, Deeks JJ, Altman DG. Measuring inconsistency in meta-analyses. *BMJ* 2003; 327:557–60. <https://doi.org/10.1136/bmj.327.7414.557> PMID: 12958120
15. Veroniki AA, Vasiliadis HS, Higgins JP, Salanti G. Evaluation of inconsistency in networks of interventions. *Int J Epidemiol*. 2013; 42(1):332–45. <https://doi.org/10.1093/ije/dys222> PMID: 23508418.

16. Chaimani A, Higgins JP, Mavridis D, Spyridonos P, Salanti G. Graphical tools for network meta-analysis in STATA. *PLoS one*. 2013; 8(10):e76654. Epub 2013/10/08. <https://doi.org/10.1371/journal.pone.0076654> PMID: 24098547; PubMed Central PMCID: PMC3789683.
17. Rucker G, Schwarzer G. Ranking treatments in frequentist network meta-analysis works without resampling methods. *BMC medical research methodology*. 2015; 15:58. Epub 2015/08/01. <https://doi.org/10.1186/s12874-015-0060-8> PMID: 26227148; PubMed Central PMCID: PMC4521472.
18. De Palma GD, di Matteo E, Romano G, Fimmano A, Rondonone G, Catanzano C. Plastic prosthesis versus expandable metal stents for palliation of inoperable esophageal thoracic carcinoma: a controlled prospective study. *Gastrointest Endosc*. 1996; 43(5):478–82. Epub 1996/05/01. PMID: 8726762.
19. Homs MY, Steyerberg EW, Eijkenboom WM, Tilanus HW, Stalpers LJ, Bartelsman JF, et al. Single-dose brachytherapy versus metal stent placement for the palliation of dysphagia from oesophageal cancer: multicentre randomised trial. *Lancet (London, England)*. 2004; 364(9444):1497–504. Epub 2004/10/27. [https://doi.org/10.1016/s0140-6736\(04\)17272-3](https://doi.org/10.1016/s0140-6736(04)17272-3) PMID: 15500894.
20. Homs MYV, Wahab PJ, Kuipers EJ, Steyerberg EW, Grool TA, Haringsma J, et al. Esophageal stents with antireflux valve for tumors of the distal esophagus and gastric cardia: a randomized trial. *Gastrointestinal Endoscopy*. 2004; 60(5):695–702. [http://dx.doi.org/10.1016/S0016-5107\(04\)02047-4](http://dx.doi.org/10.1016/S0016-5107(04)02047-4). PMID: 15557944
21. Knyrim K, Wagner HJ, Bethge N, Keymling M, Vakil N. A controlled trial of an expansile metal stent for palliation of esophageal obstruction due to inoperable cancer. *New England Journal of Medicine*. 1993; 329(18):1302–7. doi: 10.1056/NEJM199310283291803. PMID: 7692297
22. Laasch H-U, Marriott A, Wilbraham L, Tunnah S, England RE, Martin DF. Effectiveness of Open versus Antireflux Stents for Palliation of Distal Esophageal Carcinoma and Prevention of Symptomatic Gastroesophageal Reflux 1. *Radiology*. 2002; 225(2):359–65. <https://doi.org/10.1148/radiol.2252011763> PMID: 12409567
23. O'Donnell CA, Fullarton GM, Watt E, Lennon K, Murray GD, Moss JG. Randomized clinical trial comparing self-expanding metallic stents with plastic endoprostheses in the palliation of oesophageal cancer. *British Journal of Surgery*. 2002; 89(8):985–92. <https://doi.org/10.1046/j.1365-2168.2002.02152.x> PMID: 12153622
24. Power C, Byrne PJ, Lim K, Ravi N, Moore J, Fitzgerald T, et al. Superiority of anti-reflux stent compared with conventional stents in the palliative management of patients with cancer of the lower esophagus and esophago-gastric junction: results of a randomized clinical trial. *Diseases of the Esophagus*. 2007; 20(6):466–70. <https://doi.org/10.1111/j.1442-2050.2007.00696.x> PMID: 17958720
25. Sabharwal T, Gulati MS, Fotiadis N, Dourado R, Botha A, Mason R, et al. Randomised comparison of the FerX Ella antireflux stent and the ultraflex stent: Proton pump inhibitor combination for prevention of post-stent reflux in patients with esophageal carcinoma involving the esophago-gastric junction. *Journal of Gastroenterology & Hepatology*. 2008; 23(5):723–8.
26. Sabharwal T, Hamady MS, Chui S, Atkinson S, Mason R, Adam A. A randomised prospective comparison of the Flamingo Wallstent and Ultraflex stent for palliation of dysphagia associated with lower third oesophageal carcinoma. *Gut*. 2003; 52(7):922–6. PMID: 12801944
27. Shenfine J, McNamee P, Steen N, Bond J, Griffin SM. A randomized controlled clinical trial of palliative therapies for patients with inoperable esophageal cancer. *American Journal of Gastroenterology*. 2009; 104(7):1674–85. <https://doi.org/10.1038/ajg.2009.155> PMID: 19436289
28. Siersema PD, Hop WC, Dees J, Tilanus HW, van Blankenstein M. Coated self-expanding metal stents versus latex prostheses for esophagogastric cancer with special reference to prior radiation and chemotherapy: a controlled, prospective study. *Gastrointestinal endoscopy*. 1998; 47(2):113–20. PMID: 9512274
29. Siersema PD, Hop WC, Van Blankenstein M, van Tilburg AJ, Bac D-J, Homs MY, et al. A comparison of 3 types of covered metal stents for the palliation of patients with dysphagia caused by esophagogastric carcinoma: a prospective, randomized study. *Gastrointestinal endoscopy*. 2001; 54(2):145–53. PMID: 11474382
30. Vakil N, Morris AI, Marcon N, Segalin A, Peracchia A, Bethge N, et al. A prospective, randomized, controlled trial of covered expandable metal stents in the palliation of malignant esophageal obstruction at the gastroesophageal junction. *American Journal of Gastroenterology*. 2001; 96(6):1791–6. <https://doi.org/10.1111/j.1572-0241.2001.03923.x> PMID: 11419831.
31. van Heel NCM, Haringsma J, Boot H, Cats A, Vanhoutvin SALW, Kuipers EJ. Comparison of 2 expandable stents for malignant esophageal disease: a randomized controlled trial. *Gastrointestinal Endoscopy*. 2012; 76(1):52–8. <https://doi.org/10.1016/j.gie.2012.02.050> PMID: 22726466
32. Verschuur EM, Repici A, Kuipers EJ, Steyerberg EW, Siersema PD. New design esophageal stents for the palliation of dysphagia from esophageal or gastric cardia cancer: a randomized trial. *The American*

- journal of gastroenterology. 2008; 103(2):304–12. <https://doi.org/10.1111/j.1572-0241.2007.01542.x> PMID: 17900325
33. Wenger U, Johnsson E, Arnelo U, Lundell L, Lagergren J. An antireflux stent versus conventional stents for palliation of distal esophageal or cardia cancer: a randomized clinical study. *Surgical endoscopy*. 2006; 20(11):1675–80. CN-00572069 UPDATE. <https://doi.org/10.1007/s00464-006-0088-2> PMID: 16960663
 34. Wenger U, Johnsson E, Bergquist H, Nyman J, Ejnell H, Lagergren J, et al. Health economic evaluation of stent or endoluminal brachytherapy as a palliative strategy in patients with incurable cancer of the oesophagus or gastro-oesophageal junction: results of a randomized clinical trial. *European journal of gastroenterology & hepatology*. 2005; 17(12):1369–77. CN-00561518 UPDATE.
 35. White RE, Chepkwony R, Mwachiro M, Burgert SL, Enders FT, Topazian M. Randomized trial of small-diameter versus large-diameter esophageal stents for palliation of malignant esophageal obstruction. *Journal of Clinical Gastroenterology*. 2015; 49(8):660–5. <https://doi.org/10.1097/MCG.0000000000000333> PMID: 25992812
 36. Zhu HD, Guo JH, Mao AW, Lv WF, Ji JS, Wang WH, et al. Conventional stents versus stents loaded with (125)iodine seeds for the treatment of unresectable oesophageal cancer: a multicentre, randomised phase 3 trial. *The Lancet Oncology*. 2014; 15(6):612–9. CN-00988108 UPDATE. [https://doi.org/10.1016/S1470-2045\(14\)70131-7](https://doi.org/10.1016/S1470-2045(14)70131-7) PMID: 24742740
 37. Spaander MC, Baron TH, Siersema PD, Fuccio L, Schumacher B, Escorsell À, et al. Esophageal stenting for benign and malignant disease: European Society of Gastrointestinal Endoscopy (ESGE) Clinical Guideline. *Endoscopy*. 2016; 48(10):939–48. <https://doi.org/10.1055/s-0042-114210> PMID: 27626318
 38. Sgourakis G, Gockel I, Radtke A, Dedemadi G, Gourmas K, Mylona S, et al. The use of self-expanding stents in esophageal and gastroesophageal junction cancer palliation: a meta-analysis and meta-regression analysis of outcomes. *Digestive diseases and sciences*. 2010; 55(11):3018–30. Epub 2010/05/05. <https://doi.org/10.1007/s10620-010-1250-1> PMID: 20440646.
 39. Lettmaier S, Strnad V. Intraluminal brachytherapy in oesophageal cancer: defining its role and introducing the technique. *Journal of Contemporary Brachytherapy*. 2014; 6(2):236–41. doi: 10.5114/jcb.2014.43780. PMC4105652. PMID: 25097567
 40. Fuccio L, Mandolesi D, Farioli A, Hassan C, Frazzoni L, Guido A, et al. Brachytherapy for the palliation of dysphagia owing to esophageal cancer: A systematic review and meta-analysis of prospective studies. *Radiotherapy and oncology: journal of the European Society for Therapeutic Radiology and Oncology*. 2017; 122(3):332–9. Epub 2017/01/21. <https://doi.org/10.1016/j.radonc.2016.12.034> PMID: 28104297.
 41. Fuccio L, Guido A, Hassan C, Frazzoni L, Arcelli A, Farioli A, et al. Underuse of brachytherapy for the treatment of dysphagia owing to esophageal cancer. An Italian survey. *Digestive and liver disease: official journal of the Italian Society of Gastroenterology and the Italian Association for the Study of the Liver*. 2016; 48(10):1233–6. Epub 2016/08/02. <https://doi.org/10.1016/j.dld.2016.07.003> PMID: 27476465.
 42. Sabharwal T, Gulati MS, Fotiadis N, Dourado R, Botha A, Mason R, et al. Randomised comparison of the FerX Ella antireflux stent and the ultraflex stent: proton pump inhibitor combination for prevention of post-stent reflux in patients with esophageal carcinoma involving the esophago-gastric junction. *Journal of gastroenterology and hepatology*. 2008; 23(5):723–8. Epub 2008/04/16. <https://doi.org/10.1111/j.1440-1746.2008.05396.x> PMID: 18410607.
 43. Mills EJ, Ghement I, O'Regan C, Thorlund K. Estimating the power of indirect comparisons: a simulation study. *PloS one*. 2011; 6(1):e16237. Epub 2011/02/02. <https://doi.org/10.1371/journal.pone.0016237> PMID: 21283698; PubMed Central PMCID: PMC3025012.
 44. Greenland S, Mansournia MA, Altman DG. Sparse data bias: a problem hiding in plain sight. *BMJ*. 2016; 352. <https://doi.org/10.1136/bmj.i1981> PMID: 27121591
 45. Sterne JA, Gavaghan D, Egger M. Publication and related bias in meta-analysis: power of statistical tests and prevalence in the literature. *Journal of clinical epidemiology*. 2000; 53(11):1119–29. PMID: 11106885
 46. Trinquart L, Chatellier G, Ravaud P. Adjustment for reporting bias in network meta-analysis of antidepressant trials. *BMC medical research methodology*. 2012; 12(1):150. <https://doi.org/10.1186/1471-2288-12-150> PMID: 23016799